
- ()

() -

- : -
- -
: - -

:

-

(LORENZ)

(Mineral paragenesis)

5- 2,5

650 - 500

(Orthogenetic)

-

:

180

(Juteau,1980)

. (Parrot,1977)

-

-

(Parrot,1977;Parrot,1980)

:
(35,55 35,51) (35,55 35,52)
(35,55 35,50)
-
(-)
-)
(-

- ()

(Dubertret,1953) NW – SE

3 50 300 – 5
(Kazmin and . 0.5 -
- (Parrot,1980) Kulakov,1968)


.(Whitechurch and Parrot,1974) (Majer,1962) (Chenevoy,1959)
(Kazmin and Kulakov,1968) (Chenevoy,1959) (Dubertret,1953)
(Piro,1967)

. (Majer,1962)
K– Ar
(Delaloye MJ 88+3 (Thuizat etal.,1981)
(Parrot . / MJ 90 etal.,1980)
and Whitechurch,1978)

(Whitechurch and .

(Parrot,1977)

Parrot,1974)



(Parrot,1977)

هدف البحث :

()

- ()

(Lorenz,1980;1981a;1981b)

()

(Paragenic)

(Orthogenic)

طرائق التحليل :

140

Eltinor-4

Laborlux-12 Pol

) Glakov

.(

(JXA-480A)

(-)

(KEUX – EDS)

-

(Epoxi-Hart)

(Fa-Astemex-)

µa Kv

.(- :)

(XRF) X-ray fluoroescence – Spectrometry

Schmeltableten

.Lanthanium trioxide

P

K

Spektrophotometry

.(AAS) Atomic absorption spectrometry

Na

.(- :)

الدراسة البتروغرافية :

-

(Chenevoy,1959)

(Dubertret,1953)

. (Whitechurch,1977)

(Majer,1962), (Piro,1967) :

(Kazmin and Kulakov,1968)

:

-

:

-

.

-

-

:

-

-

-

.

-


()

- ()

()

- - - - - :
- :

(2 -)



- : 2 -
- Bi - Py - G - Ep - Q - Pg - Ho)
(- K.M. - Ak - O.M. - Chl
(- A - QD - BB - KE - G)

(- -)

(Lorenz,1981a;1981b)



BB-13

.(Safarjalani,1996)

B B-14

%77,6 G-12

%22,4

⋮

. 0,10 - 3,5

(ny=nz)

(nx)

(nz)

(ny)

(nx)


(3 -)

(4 -)

(-)

-

()



X A -



(+) X BB -

° 56 ° 124 (110)

° 12 ° 25-° 8 0,015 – 0,026

.Gemeine Hornblende

Fe⁺³/Fe⁺² (Leake,1978)

(Papike et al.,1974)

1.34<(Ca+Na)_x

0.67>Na_x

A (Na+K)

Al₍₄₎ Si_(z) Ti_(y) Al₍₆₎/Fe⁺³

(Na+K)_A Al₍₄₎ . (Na+K)_A

(Leake,1978)

.(Safarjalani,1996) (–) Hastingsit – Tschermakit

:

% 28,4 - % 22,1

-	()
0,60	0,85	0,20
:		:

(Safarjalani, 1997)

- An= %66-%42

: _____ .(-)

. 0,4 - 0,08

.

:

%4,3 - %2,8

1,2 0,50 -0,06



G - (-) X



(-) X G -

جدول رقم (١) التركيب الكيميائي لفتل الهورنبلند - أمفيبوليت منطقة البابر والبسيط (سورية)

Lok	G	G	G	KE	KE	KE	BB	BB	BB	QD	A	A	A	A	A	A
Pro	G6	G12	G13	KE10	KE12	KE13	BB9	BB10	QD4	A2	A8	A14	A15	A17		
SiO2	45.50	44.85	54.65	46.81	42.05	48.42	49.17	43.22	49.87	45.68	50.20	46.25	42.72	47.07		
TiO2	1.54	0.66	1.39	1.16	2.09	1.90	0.78	1.12	0.65	1.24	0.72	1.96	2.04	2.82		
Al2O3	11.97	13.56	11.65	9.75	13.08	11.30	9.78	13.23	9.89	12.27	15.25	11.18	13.39	11.65		
Fe2O3	0.70	3.51	0.00	0.77	3.86	1.93	0.72	1.04	0041	0.90	0.00	1.34	1.05	0.00		
MgO	18.79	14.58	13.20	13.14	10.71	15.30	13.18	3.64	18.29	14.37	12.97	13.90	10.77	12.43		
CaO	12.17	11.51	12.60	12.38	12.05	11.56	12.62	11.91	12.15	11.95	12.37	12.32	12.10	11.58		
FeO	11.98	9.47	12.87	13.21	10.90	6.94	13.37	17.77	7.01	10.74	7.51	10.35	14.81	11.95		
Na2O	1.72	1.60	1.82	1.21	1.49	0.94	1.11	1.49	1.60	1.99	0.86	2.15	1.89	1.44		
K2O	0.79	0.26	0.96	0.91	1.15	0.41	0.33	1.94	0.46	0.60	0.10	0.14	1.32	1.05		
Summe	100.1	100.4	100.2	100.1	100.4	100.2	100.1	100.4	100.0	100.1	100.0	100.1	100.1	100.0		
Kation=23(0)																
Si	6.50	6.35	6.55	6.72	6.14	6.71	6.95	6.38	6.90	6.49	6.87	6.58	6.24	6.68		
Al(4)	1.50	1.65	1.45	1.28	1.86	1.29	1.05	1.62	1.11	1.51	1.14	1.42	1.76	1.32		
Summe (z)	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00		
Al(6)	0.52	0.61	0.52	0.87	0.40	0.64	0.60	0.68	0.51	0.54	1.32	0.45	0.55	0.63		
Fe3	0.08	0.37	0.00	0.08	0.43	0.20	0.08	0.12	0.04	0.10	0.00	0.14	0.12	0.00		
Ti	0.16	0.07	0.15	0.17	0.23	0.20	0.08	0.12	0.07	0.13	0.07	0.18	0.22	0.30		
Mg	2.94	3.08	2.82	2.87	2.33	3.16	2.78	1.90	3.77	3.12	2.46	2.95	2.35	2.63		
Fe2	1.31	0.87	1.51	1.50	1.62	0.80	1.46	2.18	0.62	1.12	0.86	1.27	1.77	1.42		
Summe (y)	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00		
Fe	0.12	0.25	0.04	0.08	0.08	0.01	0.00	0.01	0.20	0.16	0.00	0.02	0.04	0.00		
Ca	1.86	1.75	1.94	1.90	1.98	1.72	1.92	1.88	1.80	1.82	1.18	1.88	0.89	1.76		
Al	0.02	0.01	0.02	0.02	0.03	0.27	0.08	0.11	0.00	0.02	0.19	1.10	0.07	0.24		
Summe (x)	2.00	2.01	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00		
Na	0.46	0.43	0.48	0.32	0.39	0.25	0.22	0.32	0.34	0.53	0.04	0.49	0.47	0.16		
K	0.14	0.11	0.18	0.17	0.21	0.07	0.06	0.37	0.09	0.11	0.02	0.07	0.25	0.19		
Summe (A)	0.61	0.54	0.66	0.49	0.60	0.32	0.28	0.69	0.43	0.64	0.06	0.56	0.72	0.35		

جدول رقم (٢) التركيب الكيميائي لفلز الالاجيوكلاز - أمفيبوليت منطقة البائر والبسيط (سورية)

Lok.	G																							
	G6	G12	G13	KE10	P1	P2	P3	KE12	P1	P2	KE13	P1	P1	BB13	QD4	A2	A8	A14	A15	P1	P2	A17		
PP.Nr.																								
S02	69.68	56.88	60.32	52.25	51.44	47.98	53.09	57.59	55.73	64.63	60.92	52.00	59.43	60.37	59.52	52.53	52.47	52.53	52.47	52.47	52.47	52.47	64.31	64.31
A1203	20.92	20.39	19.99	29.01	29.51	33.56	27.15	23.75	28.26	20.33	19.99	28.67	22.95	22.88	23.27	29.12	29.18	29.12	29.18	29.18	29.18	29.18	37.03	37.03
Fe203	0.00	3.43	1.98	0.00	0.00	0.00	1.79	1.84	0.00	0.00	1.98	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.39	0.39	
CaO	8.57	14.46	12.85	14.28	14.60	14.39	12.22	11.11	11.97	8.66	12.85	14.24	12.77	12.21	12.43	14.11	14.03	14.11	14.03	14.03	14.03	12.78	12.78	
Na2O	5.82	3.20	4.27	4.36	4.36	4.08	5.06	5.71	4.04	6.37	4.27	5.09	4.85	4.54	4.78	4.24	4.42	4.42	4.42	4.42	4.42	4.49	4.49	
K2O	0.00	1.63	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
Sum.	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	
Katinsw(0)																								
Si	2.91	2.63	2.75	2.38	2.35	2.20	2.43	2.61	2.50	2.86	3.75	2.38	2.67	2.70	2.67	2.39	2.39	2.39	2.39	2.39	2.39	2.42	2.42	
Al	1.04	1.11	1.06	1.56	1.59	1.81	1.46	1.27	1.50	1.06	1.06	1.06	1.22	1.21	1.32	1.56	1.57	1.56	1.57	1.57	1.57	1.46	1.46	
Fe+3	0.00	0.12	0.07	0.00	0.00	0.00	0.06	0.06	0.00	0.00	0.07	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01	
Sum.	3.95	3.85	3.88	3.94	3.94	4.01	3.95	3.94	4.00	3.92	3.88	3.93	3.89	3.91	4.00	3.95	3.95	3.95	3.95	3.95	3.95	4.07	4.07	
Ca	0.38	0.72	0.62	0.70	0.72	0.71	0.63	0.54	0.56	0.41	0.62	0.60	0.62	0.45	0.42	0.60	0.68	0.62	0.68	0.68	0.68	0.62	0.62	
Na	0.47	0.92	0.37	0.39	0.39	0.36	0.45	0.50	0.35	0.55	0.37	0.45	0.42	0.39	0.42	0.39	0.39	0.42	0.39	0.39	0.39	0.33	0.33	
K	0.00	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
Sum.	0.68	1.10	1.00	1.09	1.10	1.07	1.08	1.04	0.99	0.96	1.00	1.05	1.04	0.98	1.01	1.06	1.07	1.06	1.07	1.07	1.07	0.94	0.94	
An	44.84	65.16	62.45	63.89	64.77	66.09	65.53	61.82	62.09	42.90	62.45	60.73	59.27	59.78	58.97	46.78	36.96	46.78	36.96	36.96	36.96	61.14	61.14	
Ab	55.16	26.09	37.55	36.11	35.23	33.91	41.47	48.18	37.91	57.10	37.55	39.27	40.73	40.22	41.03	35.22	36.31	35.22	36.31	36.31	36.31	38.86	38.86	
Or	0.00	8.75	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
Int(ant/cab)	0.21	0.92	0.51	0.57	0.61	0.67	0.34	0.07	0.47	-0.29	0.51	0.28	0.38	0.40	0.36	0.61	0.56	0.61	0.56	0.56	0.56	0.63	0.63	

- ()

.
.
:
%6,2 %1,5

(ny=nz)

(nx)

0,042- 0,035

-

.
:
:

6 - 1,8

(Parrot,1977)

.(-)

: _____

(G-15,G-13,G12,G11,G-10,G-9)

(QD-1 ,QD-2)

0,48 – 0,06

.(-)

: _____

Opt.(-),l(+)

Opt.(+),l(-)

0,066-0,044

_____ - () _____

_____ . - _____
_____ : _____

0,028 -0,020 ()

_____ . _____
_____ . _____
_____ : _____

_____ . _____
_____ : _____

_____ - 3,35 - 1,20

(0112) (1011)

() ()

_____ . _____
_____ . _____

:

. - - :
% 0,8 - % 2,1 .

تحديد البارا جينيز الفلزي وشروط التحول الإقليمي :

: () -
-) + :
± ± + ()
± ()
± ± ()
.
-) + :
± ± + ()
.
± ±

- ()

() + :
± ± ± +

() + :
) ± ± +
± ± ± (

+ () + :
± ± ±

:
± ± + (-) +
%1 >

(Mathe,1985)

(Winkler,1965)

- - -
()
-
()
-
-

النتائج :

(2-) :

- -

% 28,4 % 43,8

.% 19,3 % 68,9

(Walker et al.,1960)

% 5 -

% 20 BB-13

. (Mathe,1985)

- -

(-)

-

()



(-) X QD -



(+) X BB -

:

- - :

.

-

.

()

(Bucher and Frey,1994) (Wimmenauer,1985)

± ± + + +) :

± ± +) (

.(± ±

-

QD-2 QD-1

()

.

-

(Wimmenauer,1985)

(Pichler,1993)

- ()

()

()

(Parrot,1980)

(+ + +)

- (Roesler,1983) (Seim,1967)

(Pichler,1993)

+ +)

- (

5 – 2,5

650--500

(Spear,1980;1981a; (Plyusnina,1982) (Brown,1977)

.1981b)

الدراسة الجيوكيميائية:

:()

()

()

. -
(Leake,1964)

mg c

(Leake,1964) (-)

-

100 mg - c - (al - alk) (Leake,1964)

G-15, G-13

(Walker et al.,1960)

(-) MgO-CaO- FeO*

Walker

G15,G13

(Evans and Leake,1960)

mg fm

		-		()	
(Van de Kamp,1968)					. mg
	c		al-alk		.
				(Beuge,1989)	
				(Na - Ca - Mg)	
			.Na ₂ O.(CaO) ²	(CaO) ² / (MgO) ²	
				(Beuge)	
	(Frohlich,1960)			(-)	
		Ni		Cr	
ppm- 20-150		Cr		ppm-20	Cr
			ppm- 150		Cr
Co		ppm- 40		(Lange,1965)	.
.			V		ppm- 200
		(Siegferd,1988)			. QD-6 A-13
c/fm					al-alk-c/fm
				(1,50-.25)	
			c/fm		
					. G - 13 G - 15

جدول (٣)

العناصر الكيميائية الرئيسية مقدرة بـ % الداخلة في تركيب أمفيبوليت منطقة البابر والبسيط (سورية)

Sum.	LiO	P2O5	K2O	Na2O	CaO	MgO	MnO	Fe2O3	Al2O3	TiO2	SiO2	Pro.Nr
100.1	0.82	0.25	1.3	3.32	8.56	8.94	0.17	13.12	15.9	2.62	45.1	G-01
99.98	1.80	0.15	0.95	2.75	8.87	8.71	0.17	11.33	16.81	1.32	47.12	G-02
100.75	1.11	0.12	0.72	2.71	10.16	10.67	0.18	14.55	14.48	2.45	43.60	G-03
99.83	1.24	0.34	1.80	2.50	6.70	9.45	0.18	12.80	14.66	3.16	47.00	G-04
99.65	1.68	0.32	1.18	3.51	10.50	6.98	0.15	12.81	15.06	2.68	44.78	G-05
99.66	0.80	0.25	0.73	3.97	7.72	7.63	0.13	15.34	15.45	2.34	45.30	G-06
99.92	0.65	0.35	1.50	2.75	8.08	9.90	0.17	14.13	15.14	3.15	44.10	G-07
99.88	1.66	0.32	0.93	2.39	11.58	8.92	0.16	9.78	15.12	2.72	46.30	G-08
99.91	0.68	0.25	1.14	1.70	14.90	9.48	0.15	10.66	13.58	2.17	45.20	G-09
100.07	1.55	0.31	0.51	3.09	7.90	7.88	0.13	10.49	15.24	0.72	52.25	G-10
99.96	1.12	0.35	1.00	3.16	10.40	8.07	0.18	13.10	14.76	2.62	45.20	G-11
99.92	1.59	0.35	0.48	2.91	8.23	6.77	0.13	12.24	15.12	0.93	51.17	G-12
100.05	1.72	0.23	0.50	1.16	29.73	6.27	0.22	7.71	7.93	1.68	42.90	G-13
99.27	1.83	0.19	0.08	2.58	9.58	8.42	0.16	11.80	16.09	1.46	47.08	G-14
100	1.87	0.27	0.37	0.63	32.04	3.02	0.17	8.18	9.8	2.22	41.43	G-15
100.06	1.74	0.36	0.74	4.19	8.54	7.53	0.14	12.78	14.31	2.48	47.25	G-17
99.9	1.74	0.31	0.76	4.14	8.92	7.52	0.14	12.12	14.38	2.51	47.36	G-18
101.76	1.84	0.05	2.36	2.06	11.42	9.32	0.13	11.40	14.52	0.86	47.80	KE-01
99.94	1.12	0.06	1.36	3.27	11.79	9.91	0.14	12.57	13.59	1.13	45.00	KE-02
99.92	1.45	0.17	0.55	2.98	8.65	6.43	0.16	9.98	16.45	0.65	52.45	KE-03
100.22	1.62	0.22	1.06	2.46	7.74	6.51	0.18	11.31	16.91	1.37	50.84	KE-04
100.61	0.55	0.09	0.55	2.02	14.57	9.22	0.16	10.19	15.94	1.32	46.00	KE-05
100.32	0.60	0.38	1.05	2.93	11.12	9.19	0.17	12.48	11.13	2.92	48.35	KE-06
100.93	0.67	0.30	0.88	2.17	12.18	8.73	0.11	10.19	13.73	2.27	49.70	KE-07
99.96	1.54	0.24	0.63	2.78	8.10	7.15	0.17	10.84	16.34	0.82	51.35	KE-08
100.00	1.03	0.27	1.16	2.36	8.24	7.55	0.18	11.80	16.70	1.41	49.30	KE-09
99.19	0.61	0.10	0.75	3.61	11.54	9.16	0.16	10.34	15.79	1.23	45.90	KE-10
100.36	1.20	0.40	0.82	2.32	13.31	8.90	0.18	13.75	11.04	2.74	45.70	KE-11
100.97	0.71	0.66	0.44	2.11	15.82	6.96	0.16	11.93	14.75	1.33	46.10	KE-12
99.89	0.65	0.13	0.47	2.43	14.37	8.35	0.14	12.18	15.37	1.40	44.40	KE-13
99.79	1.66	0.15	0.50	3.94	11.75	6.86	0.18	13.42	14.55	1.29	45.49	KE-14
99.07	0.78	0.25	1.14	2.60	12.87	9.64	0.17	11.24	12.97	3.51	43.90	KE-15
100.58	1.21	0.06	0.74	3.15	12.27	9.19	0.15	12.66	12.37	1.78	47.00	KE-16
98.83	1.27	0.18	0.74	2.47	12.14	11.97	0.15	11.54	11.67	1.50	45.20	KE-17
99.37	0.92	0.07	0.46	3.47	14.55	9.10	0.25	10.85	14.32	1.08	44.30	KE-18
99.90	1.40	0.32	1.13	2.78	11.36	8.44	0.20	12.51	14.26	2.90	44.60	KE-19
99.16	1.24	0.30	0.81	1.39	12.27	8.68	0.19	9.29	12.48	2.21	50.30	BB-01
99.64	0.15	0.42	0.32	1.11	13.44	7.35	0.19	10.62	15.95	3.89	46.20	BB-02
100.70	0.45	0.31	0.73	3.30	12.54	8.67	0.11	12.61	13.23	2.65	46.10	BB-03
100.64	1.44	0.24	0.59	2.14	10.08	10.04	0.16	12.79	12.46	2.70	48.00	BB-04
100.50	0.45	0.52	0.60	2.82	9.65	7.77	0.12	11.69	12.49	2.19	52.20	BB-05

-

()

تابع جدول (٣)
العناصر الكيميائية الرئيسية مقدرة بـ % الداخلة في تركيب أمفيبوليت منطقة البابر والبسيط (سورية)


Sum.	LIO	P2O5	K2O	Na2O	CaO	MgO	MnO	Fe2O3	Al2O3	TiO2	SiO2	Pro.Nr
99.73	1.26	0.47	0.59	3.82	10.43	6.56	0.12	12.22	13.79	3.37	47.10	BB-06
100.18	1.20	0.39	0.93	3.06	12.00	10.64	0.14	11.39	12.61	3.32	44.50	BB-07
99.63	0.67	0.41	0.92	1.17	11.69	7.74	0.14	14.09	12.78	2.12	47.90	BB-08
99.95	1.62	0.72	0.77	3.29	10.83	8.97	0.12	10.83	12.87	3.10	46.83	BB-09
98.84	0.61	0.29	0.15	1.02	11.49	10.97	0.14	11.88	12.80	2.29	47.20	BB-10
99.89	3.03	0.32	0.14	2.84	10.95	8.33	0.13	11.34	15.17	1.61	46.03	BB-11
98.69	1.36	0.21	0.60	4.47	9.68	8.14	0.18	11.28	14.20	1.67	46.90	BB-12
99.93	1.06	0.45	0.46	3.49	11.13	6.80	0.12	12.29	14.42	3.15	46.56	BB-14
98.78	1.39	0.47	0.50	3.66	9.58	8.87	0.13	11.89	13.75	2.54	46.00	QD-01
100.32	0.84	0.22	0.70	1.47	12.21	7.13	0.16	11.65	12.32	1.12	52.50	QD-02
98.86	0.87	0.05	0.28	2.13	13.06	10.05	0.19	9.37	13.31	0.95	48.60	QD-03
100.00	0.98	0.03	0.70	1.40	12.60	11.40	0.18	10.60	14.73	0.88	46.50	QD-04
98.79	1.12	0.02	0.86	1.93	12.55	11.49	0.15	9.31	14.42	0.74	46.20	QD-05
100.13	3.13	0.02	1.57	2.07	10.41	6.74	1.11	11.76	15.30	0.36	47.66	QD-06
100.00	1.24	0.35	1.98	2.06	10.32	8.34	0.21	12.22	16.12	3.63	43.53	QD-07
101.01	1.30	0.43	1.84	2.55	11.35	8.37	0.21	12.08	15.20	3.59	44.09	QD-08
99.66	0.87	0.34	0.19	2.67	11.93	7.95	0.17	14.72	13.73	1.99	45.10	A-01
100.46	0.67	0.07	0.51	2.58	13.78	9.71	0.16	9.76	14.03	0.89	48.30	A-02
100.21	0.56	0.10	0.67	1.49	14.63	7.38	0.20	12.15	15.76	2.07	45.20	A-03
101.32	0.68	0.05	0.75	2.56	13.59	8.90	0.15	11.41	15.48	1.25	46.50	A-04
100.18	1.01	0.06	0.48	3.17	11.46	9.52	0.15	10.17	15.34	1.32	47.50	A-05
100.73	0.72	0.07	0.66	2.51	14.43	7.31	0.09	12.14	14.65	0.95	47.20	A-06
100.44	0.89	0.08	1.07	2.90	11.27	8.48	0.19	11.55	14.27	0.84	48.90	A-07
99.82	0.71	0.08	0.37	3.47	12.77	8.25	0.16	11.12	13.12	1.57	48.20	A-08
99.43	0.95	0.06	0.20	2.12	10.98	11.21	0.18	11.93	13.63	1.27	46.90	A-09
100.79	0.67	0.06	0.20	2.20	11.85	9.17	0.21	10.95	14.27	0.91	50.30	A-10
100.50	0.87	0.08	0.50	2.45	12.12	8.39	0.18	11.08	14.50	1.53	48.80	A-11
100.42	0.76	0.09	0.24	3.29	13.25	7.15	0.18	10.45	14.04	0.97	50.00	A-12
100.55	1.93	0.18	0.66	2.85	9.12	6.95	0.18	10.08	16.42	0.62	51.56	A-13
99.94	1.76	0.21	0.63	2.61	8.87	7.13	0.17	11.13	16.31	0.68	50.44	A-14
98.95	1.19	0.10	0.71	2.20	13.30	9.06	0.15	10.01	15.60	1.33	45.30	A-15
98.74	1.08	0.06	0.80	3.07	14.60	10.06	0.13	9.51	14.23	1.00	44.20	A-16
100.00	1.86	0.18	1.30	3.52	10.06	10.18	0.14	9.32	14.41	1.77	47.26	A-17
100.08	1.39	0.05	0.45	2.64	11.23	8.21	0.16	9.13	15.08	0.91	50.83	A-18

جدول (٤)
العناصر الكيميائية الثانوية مقدرة بـ Ppm الداخلة في تركيب أمفيبوليت منطقة البايير والبسيط (سورية)



V	Co	Cr	Ni	Zn	Rb	Sr	Y	Zr	Pb	Cu	Pro.Nr
		165	112	122	29	698	26	105	17	119	G-01
249	83	204	118	103	36	934	40	143	19	93	G-02
		234	196	129	13	152	39	169	24	70	G-03
		135	126	94	16	324	30	123	32	99	G-04
222	70	195	109	124	27	626	31	148	26	127	G-05
		185	102	77	20	690	31	144	16	74	G-06
		146	126	104	67	275	30	148	17	130	G-07
		99	74	116	26	295	47	210	19	131	G-08
		122	80	83	22	419	25	141	27	139	G-09
254	68	149	118	84	48	876	40	149	20	156	G-10
		194	119	117	18	252	50	190	20	116	G-11
221	56	149	113	97	37	785	34	235	36	148	G-12
		290	108	86	< 10	332	26	114	20	139	G-13
204	65	174	100	105	32	678	33	149	21	105	G-14
		88	91	71	< 10	672	25	124	35	105	G-15
		238	125	85	12	659	28	120	18	105	G-17
234	67	210	124	98	< 10	706	34	134	18	89	G-18
225	69	588	190	61	< 10	490	26	102	24	80	KE-01
		344	190	75	22	272	30	119	14	78	KE-02
224	44	153	132	83	< 10	820	39	156	19	119	KE-03
266	43	230	170	65	< 10	312	30	122	17	100	KE-04
		537	143	87	15	699	31	147	16	105	KE-05
		194	118	108	< 10	200	48	278	21	150	KE-06
		241	125	90	< 10	212	48	256	20	122	KE-07
241	43	191	119	93	< 10	711	43	178	18	117	KE-08
280	45	176	134	76	< 10	249	26	131	17	115	KE-09
		191	138	86	< 10	447	20	94	16	134	KE-10
		241	114	110	< 10	722	45	290	19	129	KE-11
		162	104	68	< 10	280	28	128	14	111	KE-12
		215	168	59	< 10	435	36	177	< 18	100	KE-13
357	66	139	85	100	< 10	545	53	112	20	116	KE-14
		210	143	51	< 10	189	53	181	46	69	KE-15
		246	163	71	12	633	36	100	27	70	KE-16
		196	285	85	< 10	152	43	116	26	100	KE-17
		181	132	68	< 10	291	19	108	25	91	KE-18
		246	68	104	16	842	34	245	23	143	KE-19
		462	446	55	< 10	693	33	229	< 10	68	BB-01
		250	157	46	< 10	1777	60	365	33	119	BB-02
		156	109	70	16	421	37	267	25	116	BB-03
		505	265	116	< 10	254	37	224	18	112	BB-04
		196	131	103	18	971	40	207	16	114	BB-05

تابع جدول (٤)
العناصر الكيميائية الثانوية مقدرة بـ PPM الداخلة في تركيب أمفيبوليت منطقة البايير والبسيط (سورية)

V	Co	Cr	Ni	Zn	Rb	Sr	Y	Zr	Pb	Cu	Pro.Nr
305	78	174	98	97	< 10	517	51	248	20	89	BB-06
		304	216	80	< 10	711	23	214	< 10	106	BB-07
		238	183	102	12	571	37	254	16	135	BB-08
255	73	351	217	100	< 10	432	39	274	17	90	BB-09
		363	395	90	< 10	130	26	170	25	191	BB-10
233	92	376	454	74	< 10	116	29	152	16	95	BB-11
		269	221	72	< 10	807	40	350	< 10	140	BB-12
216	83	120	106	86	< 10	619	31	224	17	170	BB-14
		215	143	70	< 10	554	25	151	24	79	QD-01
		747	573	62	< 10	212	36	125	23	73	QD-02
		271	231	78	< 10	225	30	210	22	76	QD-03
		318	239	31	< 10	579	17	210	22	74	QD-04
123	46	454	228	100	< 10	457	42	289	22	117	QD-05
		693	280	59	45	462	20	210	12	61	QD-06
		448	282	87	34	1061	40	331	< 10	61	QD-07
		358	290	79	31	925	36	218	< 10	53	QD-08
		147	131	109	27	312	23	131	27	99	A-01
		315	154	56	< 10	180	42	139	22	76	A-02
		249	174	74	< 10	87	28	143	25	101	A-03
		272	130	66	13	322	64	143	21	101	A-04
		307	98	62	< 10	722	31	86	38	66	A-05
		349	143	76	15	309	25	98	23	98	A-06
		426	184	57	20	396	17	109	21	62	A-07
		326	104	76	< 10	370	30	116	22	74	A-08
226	38	259	99	90	< 10	317	31	124	16	94	A-09
		299	129	57	< 10	249	28	128	12	75	A-10
		234	109	62	< 10	492	29	154	< 10	73	A-11
		146	139	75	< 10	462	28	180	< 10	64	A-12
		272	187	78	< 10	272	29	131	25	92	A-13
		243	41	219	134	45	< 10	454	29	135	< 10
256	188	71		< 10	232	34	143	< 10	69	A-15	
254	185	76		13	451	23	128	14	96	A-16	
273	158	50		15	420	23	131	15	77	A-17	
		301	193	62	< 10	247	23	146	21	69	A-18




(Leake, 1964)



(Walker et al., 1960)

MgO-FeO* : -



. (Beuge, 1989) $Na_2O \cdot (CaO)^2$ $(CaO)^2 / (MgO)^2$: -

G - 15 , G - 13

تحديد طبيعة الأصل المهلي للصخور المدروسة :

(Miyashiro,1975a;1975b)

(K, Na)

(Vallance,1974) CaO Na₂O

(La Roche etal.,1980)

$$R1 = 4Si - 11 (Na + K) - 2 (Fe + Ti) \quad ; \quad R2 = 6 Ca + 2 Mg + Al$$

R1 R2

SiO₂ Na₂O+K₂O (Middlemost,1980)

K₂O+Na₂O (Irvin and Baragar,1971)

SiO₂

- ()

(Irvin and Baragar,1971)

ICPW

)

.(subalkalibasalt

P₂O₅

(Winchester and Floyd,1976)

Zr

(Miyashiro and Shido,1975)

FeO* / MgO

TiO₂

FeO* / MgO

FeO*

FeO* / MgO

TiO₂

FeO* / MgO

FeO*

FeO*

SiO₂

(Miyashiro and Shido,1975)

Cr

SiO₂

MgO/

Al_2O_3 Na₂O + K₂O- FeO-MgO (Irvin and Baragar,1971)

. (NPC)

Al_2O_3 - (FeO + TiO₂) - MgO (Jensen,1976)

(Jensen,1976)

تحديد المواقع الجيوتكتوني للصخور المهليه الأم:

:

(Pearce and Gale,1977) , (Beccaluva etal.,1979), Ti,P,Mn,Zr,Nb, Y,Cr

.(Pearce,1975) , (Rittmann,1973)

- ()

(Pearce and Gale,1977)

Ti/Y Zr/Y (Zr , Ti , Y)

F3 , F2 , F1

F1 (Pearce et al.,1976)

(W.P.B) (-) F2

(O.F.B)

(SHO) (CAB+LKTb)

(-) F3 F2

(LKTb) (LAB)

F2 F1

(W.P.B)

(O.F.B)

Ni Ti / Cr

(Beccaluva et al.,1979)

(Rittmann,1973)

.

(Pearce and Cann,1971)

(MORB)

Ba , K,P, Sr , Zn


.Nb, Cr, Y, Ti, Zr

Zr , Sr, P , K ,Rb


(Huch,1988)

(-) Cr , Y


.(Kosters,1991)



$F1 = 0.0088SiO_2 - 0.0774TiO_2 + 0.0102Al_2O_3 + 0.0066FeO - 0.0017MgO - 0.0143CaO - 0.0155Na_2O - 0.0007K_2O$
 $F2 = -0.013SiO_2 - 0.0185TiO_2 + 0.0102Al_2O_3 - 0.0134FeO - 0.03MgO - 0.0204CaO - 0.0481Na_2O + 0.0715K_2O$
 $F3 = -0.0221SiO_2 - 0.0532TiO_2 - 0.0361Al_2O_3 - 0.0016FeO - 0.031MgO - 0.0237CaO - 0.0614Na_2O - 0.0289K_2O$
 شكل - ١٢ : F1 مقابل F2 عن الباحث (Pearce et al., 1976).



(Pearce et al., 1976) F3 F2 : -



(Huch,1988)

MORB

نتائج الدراسة الجيوكيميائية:

.G15, G13

-
-
-
-
-

REFERENCES:

1. Beccaluva, L.;D. Ohnenstetter and M.Ohnenstetter. (1979): Geochemical discrimination between ocean floor and island-arc tholeiites-application to some Ophiolites. In: Can. J. earth Sc.-Ottawa 16.S. 1874-1882.
 2. Beuge, P. (1989): Zur Geochemie pelitischer Gesteine im prozess der Regionalmetamorphose. Freiberg. Bergakad., Diss. B.
 3. Brown. E.H., (1977) : The crossite content of Ca-amphibole as a guide to pressure of metamorphism. - Journ. Petrol.. 18:53-72.
 4. Bucher.K. and M.Frey., (1994):Petrogenesis of Metamorphic Rocks.,Springer Verlag,318 S.
 5. Chenevoy.M.,(1959) : le substratum metamorphique des roches vertes dans le Baer et le Bassit. NOtes et Memoires sur le Moyen-Orient,T.7.1-18.
 6. Delaloye, M.; De H.Souza; J.J. WagnerandI.Hedley (1980b): Isotopic ages on ophiolites from the eastern Mediterranean. In: A.Panayiotou (Ed.), Ophiolites. Proc. Intern. Ophiolite Symp. , Cyprus 1979, P.292-295.
 7. Dubertret, L. (1953): Geologie des roches vertes du nordouest de la Syria et du Hatay (Turquie). Notes Mem. Moyen-Orient. 6.227p.
 8. Evans, B.W.andB.E. Leake (1960): The composition and origin of the striped amphibolites of Connemara, Ireland. - In: J.Petrol. - Oxford 1., p.337-363.
 9. Frohlich, F. (1960): Ein Beitrag zur Geochemie des chroms. Geochem. Cosm. Acta 20, 215-240.
 10. Huch, K.M. (1988): Die Panafrikanische Khzama-Geosuture im zentralen Anti-Atlas. Marakko -172S. ; Berlin (SchelzkyandJeep).
 11. Irvin, T.N.andW.R.A. Baragar. (1971): A guide to the classification of the common volcanic rocks. - In: Can J.Earth Sc. Ottawa 8. p. 523-548.
 12. Jensen, L.S. (1976): A newcation plot for classifying subalkaline volcanic rocks. - In: Ontario Dep. Mines: Paper. - Ontario, 66.
 13. Juteau, T. (1980): Ophiolites of Turkey. - Ofioliti (Spec. Issue) 2:199-237p.
 14. Kazmin, V.G,andV.V Kulakov, (1968): Geological Map of Syria sheet Bassit 1/50,000 explanatory notes: 124p. Mini. of petrol. and Mineral Resources,Syria.
-

15. Kesters, B. (1991): Geochemie und Petrogenese des Peridotit und der Chromitervorkommen des Ophiolith-Komplexes im Bassit Gebiet (NW-Syrien). Diss. Naturwiss. Munster, 125S.
 16. Lange, H. (1965): Zur Genese der Metabasite im sachsichen Erzgebirge. - Leipzig, 136 S. - (Freiberger Forschungshefte: C-177).
 17. La Roche, H. De.; J.Leterrier; P. Grandclaude; M.Marchal. (1980): A classification of volcanic and plutonic rocks using R1-R2- diagram and major-element analysis. - In: Chem. Geol. - Amsterdam 29., P. 721-736.
 18. Leake, B.E. (1978): Nomenclature of amphibolites. - Amer. Mineral., 63: 1023-1052, Washington D.C.
 19. Leake, B.E. (1964):The chemical distinction between Ortho-and Paraamphibolites.-In:J.Petrol.-Oxford 5.p.238-254
 20. Lorenz. W. ,(1980) : Petrographische Nomenklatur metamorpher Gesteine. I. Grundlagen. Konzeption. Systematik. In: Z.geol. Wiss. - Berlin 8;12. - S.1479- 1509.
 21. Lorenz. W., (1981a) : Petrographische Nomenklatur metamorpher Gesteine. II. Klassifikatorische und Nomenklatorisch Spezialfragen. In: Z. Geol. Wiss. - Berlin 9 - S. 137-156.
 22. Lorenz. W., (1981b) : Petrographische Nomenklatur metamorpher Gesteine. III. Generalisierung und Extrapolationen. In:Z. Geol. Wiss. - Berlin 9-4. - S.417-426.
 23. Majer,V., (1962) :Magmatisch Gesteine um Gebiet von Bassit zwischen Latakia und Kessab im nordwestlichen Syrien.Neves.Miner.,Abh.,98.250-282S.
 24. Mathe.G.,(1985):Metamorphite.In:Pfeiffer,L;Kurze,M;Mathe,G:Einführung in die Petrologie.-Berlin.442-565S.
 25. Middlemost, E.A.K. (1980): A contribution to the nomenclature and classification of volcanic rocks. - In: Geol. mag. - London. 117., p. 51-57.
 26. Miyashiro, A. (1975a): Classification, characteristics, and origin of ophiolites. - In: geol. - Chicago 83. p. 249-281.
 27. Miyashiro, A. (1975b): Volcanic rock series and tectonic setting. - In: Ann. rev. earth plan. Sc. - palo. Alto 3., p. 251-269.
-
-

28. Miyashiro, A. and F. Shido. (1975): Tholeiitic and calc-alkali series in relation to the behaviors of titanium, vanadium, chromium, and nickel. In: Amer. J. Sc. - New Haven 275. p. 265-277.
29. Papike . J.J., Cameron. K.L. and Baldwin. K. (1974) : Amphibol and Pyroxenes: character of other than quadrilateral components and estimates of ferric iron from microprobe data.. Geol. Soc. Am.. Abstr. with program. 6. Boulder. Colorado.
30. Parrot, J.F. (1977): Assemblage ophiolitique du Baer-Bassit et termes effusifs du volcano - sedimentaire Tra. et Doc. de l O.R.S.T.M. (these) 333p.
31. Parrot, J.F. (1980): The Baer-Bassit (Northwestern; Syria) ophiolitic area-Ofioliti (Spec. Issue) 2, 279-296; Bologna.
32. Parrot, J.F. and H. Whitechurch (1978): Subductions anterieures au charriage nord-sud de la croûte tethysaenne: Facteur de metamorphisme du volcano-sedimentaire oceanique en schistes verts et amphibolites lies aux assemblages ophiolitiques syro-turcs. Rev. Geogr. Phys. Geol. dyn., 2:153-170.
33. Pearce, J.A. (1975): Basalt geochemistry used to investigate past tectonic environment of Cyprus., - In: Tectonophys. - Amsterdam 17. p. 15-43.
34. Pearce, J.A. and J.R. Cann (1971): Ophiolite origin investigated by discriminant analysis using Ti, Zr, and Y. - In: Earth plane. Sclett. - Amsterdam 12.p. 339-349.
35. Pearce, J.A. and G.H. Gale (1977): Identification of ore depositions environment from trace elements geochemistry associated to igneous hostrocks. - In: Volcanic processes in ore genesis/ed. M.J. JONES. London, p. 14-24. (Inst. min. metall. Geol. Geol. Sec. : spec. publ.:7).
36. Pearce, J.A.T.H. Gorman and T.C. Birkett (1976): The relationship between major element chemistry and tectonic environment of basic and intermediate volcanic rocks. - In: Earth plan. Sc. lett. - Amsterdam 36. p. 121-132.
37. Pichler. H., and Schmitt - Riegraf (1993) : Gesteinbildende Minerale im Dunnschliff. Enke. Stuttgart S.230.
38. Piro. Y., (1967) : Contribution al etude des Roches verts du Nord-Ouest de la Syrie thesis 3eme cycle Montpellier 142p. (Multigr).
39. Plyusnina. L.B., (1982) : Geothermometry and Geobarometry of Plagioclas - Hornblende bearing assemblages. - Contrib. Mineral. Petrol.. 80: 140-146 Berlin - Heidelberg 1982 .

40. Rittman, A., (1973): Stable mineral assemblages of igneous rocks. - Berlin, Heidelberg. - p. 262.
 41. Rosler. H. J., (1983) : Lehrbuch der Mineralogie VEB-Deut. Verlag fur Grundstoffindustrie.Leipzig. 833 S .
 42. Safarjalani.A., (1996) :Petrographical and Geochemical Study of Metabasitic and Amphibolitic Metamorphic Rocks in the Ophiolithe complex in the Bear and Bassit (N.W.Syria) Report On Scintific Study Department of Geology and Nuclear Ores (AECS) 110p.
 43. Safarjalani. A., and J. Eidam (1997) : Geochemistry and initial geotectonic setting of amphibolitic rocks outcropped in the Baer-Bassit area (Northwest Syria). Z.Geol.Wess. ,Band 25, Heft 5 - 6, 541 – 554 S.
 44. Seim. R., (1967) : Der Para - Biotit - Plgioklasgneis des Kyffhauserkristallins seine metatektischen Derivate und seine Einlagerungen. - Geol. 16. Beih. 56:1-85.
 45. Siegfert, M. (1988): Mineralogie:Eine Einfuhrung in die speziellen Mineralogie, Petrologie und Lagerstättenkunde. 2. Auflage.
 46. Spear. F.S., (1980) : NaSi-CaSi exchange equiliberium between plgioclase and amphibole. - Contr. Mineral. Petrol.. 72. 33-41.
 47. Spear. F., (1981a) : An experimental study of hornblende stability and compositional variability in amphibolite.. - Amer. - J.S.Sci.. 281: p. 697-734.
 48. Spear. F., (1981b) : Amphibole-plagioclase equilibria: An emprical model for the relation albite -tremolite=edenite+4quartz. -Contrib. Mineral. Petrol. 77:p.355-364.
 49. Thuizat ,R.;H.Whitechurch.,R.MontignyandT.Jutea.(1981):K-Ar dating of some infra-ophiolitic metamorphicsoles from the Eastern Meditteranean:New evidence for oceanic thrustings before obduction.Earth Planet.Sci.Lett52,302-310.
 50. Vallance, T.G., (1974): Spilite degradation of a tholeiitic basalt. - In: J.petro. - Oxford 15. -p. 79-96.
 51. Van de Kamp, P.C. (1968): Geochemistry and origin of metasediments in the Haliburton-Modac area, Southwestern Ontario. - In: Can. J. earth Sc. - Ottawa 5. p. 1337-1372.
-
-

-
-
52. Walker, K.R.; Joplin, G.A.; Lovering, J.F.; R Green (1960): Metamorphic and metasomatic convergence of basic igneous rocks stern Queensland. - In: J.Geol. Soc. Australia. - Sydney 6. -p. 149-178.
 53. Whitechurch. H., and J.F. Parrot., (1974) : Les ecailles metamorphiques Infraperidotitiques du Bear - Bassit. Cah. Orstom 6. 173-184.
 54. Whitechurch. H., (1977): Les roches metamorphiques infraperidotitiques du Baer-Bassit (Nw Syrien). Termoins de L ecaillage intraoceanique tethysien. Etude petrologique et structural. Thesis (unpubl.) Nancy. Univ. Franc. 194pp.
 55. Wimmenauer. W., (1985) : Petrographie der magmatischen und metamorphen Gesteine. Enke Verlag. - Stutt - gart. 382S.
 56. Winchester, J. A; P.A. floyd (1976): Geochemical magma type discrimination appl. to altered and metamorphosed basic igneous rocks. - In: Earth plan. Ac. lett. - Amsterdam 28. -p. 459-469.
 57. Winkler. H.G.F., (1965) : Die Genese der metamorphen Gesteine - Berlin – Heidelberg:218S.
-
-

Petrography and Geochemistry of the metamorphic amphibolitic rocks associating the Baer-Bassit ophiolithe (Northwest Syria)

A.M. Safarjalani

King Faisal University-Faculty of Food and Agriculture P.O.Box 420-31982 Al -Hofuf-Saudi Arabia.
Damascus University- Faculty of Agriculture Land. Soil reclamation Division- Damascus P.O. Box 30621- Syria.

Abstract

Petrography of metamorphic rock samples from amphibolitic rocks outcropped in Baer-Bassit area enabled the mineralogical composition, structures, textures and hence the classification of these rocks to be defined .

The plotting of minerals percentages of the studied samples onto classification diagrams of (LORENZ) confirms that all of these samples lie in the field assigned to amphibolitic rocks .

The mineralogical paragenesis of the studied samples is in conformity with those of green schist up to amphibolite facies . The mineralogical analysis carried out by the use of electron microscope on plagioclase and hornblende in amphibolite samples set a metamorphism temperature between 500 - 650 °C and a metamorphism pressure between 2,5 - 5 K bar.

The geochemistry of these samples proves clearly and decisively that these rocks are derived from initial matter of orthogenic nature, possessing the characteristics of a fresh non-altered basaltic tholeiitic composition in part hence forming an oceanic floor and partially an island arc.
