

The Effect of Induced Hypothyroidism on Fertility of Goats During Summer Season

Khaled A. AL-Busadah

Department of Physiology, Biochemistry and Pharmacology,
College of Veterinary Medicine and Animal Resources, King Faisal University
Al-Hassa, Kingdom of Saudi Arabia

Abstract :

The effects of Carbimazole (CZ)-induced hypothyroidism were investigated in goats during summer season. CZ treatment at a daily dose of 3mg/kg body weight for 11 days inhibited ovulation and corpus luteum formation. This suggested that heat stress, which usually causes a decline in thyroid hormones, may produce its effect on fertility in a similar manner to CZ-induced hypothyroidism.

Introduction:

Summer heat stress reduces fertility during summer in animals (Monty and Wolf 1974). Numerous reports have indicated that hot summer season causes a lowering of thyroid activity, resulting in low plasma concentration of thyroid hormones in most studied animals (Johnson and Vanjmack 1976; Hart *et. al*, 1979; Bell *et. al* 1985; Nixon *et. al* 1988; El Nouty *et. al* 1990). Thyroid hormones modulate metabolic processes (Ganong 2001; Browning *et. al* 1998). These metabolic hormones are involved in thermoregulation, because metabolism is a primary intrinsic source of heat (Finch 1986). This study was conducted to investigate the effect of induced hypothyroidism on fertility of goats during summer season.

Materials and Methods:

Animals: sixteen mature (2 -3 years) females Arady goats (weighing 25-30 kg) were used. They were housed in individual pens during summer season (June - August). After completion of one normal oestrous cycle (19 -20 days), the goats were randomly divided into two groups A&B. The treatment administered to each group A&B was as follows:

Animals in Group A (N = 8, Controls) received drench of saline from the Day of oestrous to Day 10 of the oestrous cycle (oestrous = Day 0).

Animals in Group B (N = 8) were treated exactly as Group A but with carbimazole (CZ) tablets (Neomercazole, Nicholas laboratories Ltd. UK). The tablets were powdered and dissolved in distilled water and given by drench at the rate of 3 mg/kg body weight.

In Groups A and B jugular blood (5ml) was collected daily by venipuncture using 23- gauge needle between Days 0 -20 of the oestrous cycle and the subsequent oestrous cycle. All blood sample were collected into heparinized nylon syringes and centrifuged at 1500 g for 10 minutes. Plasma samples were stored at -20 °C until analyzed for hormones. All female goats were checked for oestrous twice daily by using a fertile buck and considered to be in oestrous when they stood for mounting.

Hormone Analysis: Progesterone and 13.14 dihydro-15 ketoprostaglandin, F2a (PGFM), the metabolite of prostaglandin F2a were measured by specific radioimmunoassay previously validated (Homeida and Cooke, 1982; Homeida, 1986). The intra- and inter-assay coefficients of variations were 4.4% (n = 15) and 12% (n = 20) respectively, for progesterone and 5.6% (n =10) and 11 % (n =15) respectively, for PGFM. The sensitivity of the assay was 48 pg/tube for progesterone and 10 pg/tube for PGFM. Extraction efficiency was 85.4 ± 4.8 % for progesterone and 87.1 ± 5.1 % for PGFM and the results were corrected for extraction losses.

Statistical analysis: the results were compared by Student's t- test.

Results:

Similar pattern of hormone concentration during the first oestrous cycle in both Group A and B (Table 1) was observed. CZ produced no effect on luteal regression or oestrous. In the subsequent oestrous cycle, complete failure of luteal phase occurred in Group B. Progesterone values remained less than 0.1 ng/ml, which were significantly ($P < 0.001$) lower than values in Group A during days 5,10, and 15 of oestrous cycle. PGFM values were similar in the two groups, but significantly different ($P < 0.001$) on Day 15 of second oestrous cycle.

Table (1)

Mean (\pm SD) plasma concentrations of progesterone and 13,14-dihydro-15-ketoprostaglandin F2a (PGFM) during two oestrous cycles in goats treated with saline (Group A) or carbimazole (Group B).

Days of Oestrous cycle	Progesterone (ng/ml)		PGFM (pg/ml)	
	Group A	Group A	Group B	Group B
0	< 0.1	< 0.1	800 \pm 25	850 \pm 25
5	2.9 \pm 0.1	2.6 \pm 0.1	50 \pm 10	7 \pm 10
10	4.3 \pm 0.2	4.1 \pm 0.2	80 \pm 10	100 \pm 10
15	2.1 \pm 0.1	2.5 \pm 0.1	680 \pm 20	720 \pm 20
0	< 0.1	< 0.1	950 \pm 25	1050 \pm 25
50	2.8 \pm 0.1	< 0.1*	70 \pm 10	110 \pm 10
10	4.4 \pm 0.2	< 0.1*	105 \pm 10	75 \pm 10
15	2.5 \pm 0.1	< 0.1*	660 \pm 20	70 \pm 10*

* P<0.001

Discussions:

Induction of altered thyroid function by CZ in goats and laboratory animals have been reported earlier (Baquer *et al.*, 1976); Ibrahim *et al.*, 1984). The relative longer life span of goats when compared to laboratory animals, makes them of practical consideration to be used as a model to study chronic complications during altered thyroid function. The main objective of this study was to examine the role of hypothyroidism on fertility of goats during summer season when ambient temperature in this part of the world reaches a quiet high values. The results have indicated that CZ-induced hypothyroidism produced no effect on reproductive hormones initially, when administered during oestrous cycle. But in the subsequent oestrous cycle, ovulation did not occur as indicated by failure of corpus luteum formation and absence of progesterone rise thereafter. Similar results due to thyroidectomy or suppression of thyroid function in female goats have been shown to impair or even abolish normal ovarian function (Reddy *et al.*, 1990; Walkden-Brown *et al.*, 1990). Such results may be explained by the possibilities that impaired thyroid function may affect steroid feedback responses required for generation of pre-ovulatory

lutinizing hormone surge and suppression of ovulation (Webster *et al.*, 1991; Anderson and Barrell, 1998). The possibility for secretions of thyroid gland other than its hormones required for normal ovarian function cannot be ruled out.

Thyroid function in animals should decline as an acclimation response to increased heat after a few days to help alleviate heat stress (Pratt and Wettemann, 1986). This decline in thyroid function during heat stress may be due to effect of heat on hypothalamic-pituitary axis to cause reduction in thyrotropin releasing hormone which enables the animal to reduce basal metabolism (Johnson, 1987). Such decline in thyroid function may sometimes reduce fertility in animals as in case of CZ- induced hypothyroidism.

References

1. Anderson G.M and Barrell G. K (1998). Effects of thyroidectomy and thyroxine replacement on seasonal reproduction in the red deer hind. *Journal of Reproduction and Fertility* 113,239-250
2. Baquer N. Z, Cascales M, Maclean B., and Greenbaum A. (1976). Effects of thyroid hormone deficiency on the distribution of hepatic metabolites and control pathways of carbohydrate metabolism in liver and adipose tissue of the rat. *Journal of Biochemistry* 68, 403-413
3. Bell B. A, Hainen W. and Johnson H. D. (1985). Environmental heat effects on tropically evolved African pygmy goats. 17th Conference on biometeorology and aerobiology. American Meteorology Society. Boston, USA.
4. Browning R., Leiter-Browning M.L., Smith H. M and Wakefield T. (1998). Effects of ergotamine on plasma concentrations of thyroid hormones and cortisol in cattle. *Journal of Animal Science* 76, 1644-1650.
5. EL-Nouty F. D., Al-Haidary A.A. and Salah M. S. (1990). Seasonal effects on body temperature, thyroid function, blood glucose and milk production in lactating and dry Holstein cows in semi-arid environment. *Arab Gulf Journal for Scientific research*, 8, 89-103
6. Ganong W. F. (2001). *Review of Medical Physiology* 20th Ed. Lange medical books/McGraw-Hill. New York, USA.
7. Finch V. A. (1986). Body temperature in beef cattle. Its control and relevance to production in the tropics. *Journal of Animal Science* 69,2108-2114

8. Hart I. C., Nines J. A. and Morant S. V (1979). Endocrine control of energy metabolism in the cows: correlation of hormones and metabolites in high and low yielding cows for stages of lactation. *Journal of Dairy science* 62, 270-277
9. Homeida A. M. and Cooke R. G. (1982). Peripheral plasma concentration of 13, 14- dihydro-15-ketoprostaglandin F2a and progesterone around luteolysis and during early pregnancy in the goats. *Prostaglandins* 24, 313-321
10. Homeida A. M. (1986). Use of spironolactone to investigate the role of testosterone secretion during luteolysis in the goat. *Journal of Reproduction and Fertility* 76, 153-157
11. Ibrahim R. E., Maglad M. A., Adam S.E. I. Mirgany T. E. and Wasfi I. A. (1984). The effect of altered thyroid status on lipid metabolism in the Nubian goats. *Comparative Biochemistry and Physiology* 77B, 507-512
12. Johnson H. D. and Vanjmack W. J. (1976). Symposium: Sterss and Health of the dairy cow. *Journal of Dairy Science* 89, 1601617
13. Johnson H. D. (1987). Bioclimate and livestock, In: Johnson H. D.(ed.) *Bioclimatology and adaptation of Livestock*. Elsevier Science Publishers B. V. Amesterdam. The Netherlands pp. 3-16
14. Monty D. E. and Wolf I. K. (1974). Summer heat stress and reduced fertility in the Holstein-Friesian cows in Arizona. *American Journal of Veterinary Research* 35, 1495-1500
15. Nixon D. A., Akasha M. A., and Anderson C. R. (1988). Free and total thyroid hormones in serum of Holstein cows. *Journal of Dairy Science* 71, 1152.-1160
16. Pratt B. R. and Wettemann (1986). The effect of environmental temperature on concentration of thyroxine and tri-iodothyronine after thyrotropin releasing hormone in steers. *Journal of Animal Science* 62, 1346-1352
17. Reddy I. I. Varshney V. P ., Sanwalt P. C., Agarwal N and Pande J. K (1990).Peripheral plasma estradiol 17/3 and progesterone levels in female goats induced hypothyroidism. *Small Ruminant Research* 22, 149-154
18. Walkden-Brown S.W. Davidson R. H., Milton J. T. B., and Martin G. B (1990). Thyroidectomy late in the breeding season and ovulation in Cashmere goats. 13th temational Congress on Animal Reproduction Abstract pp 1-19
19. Webster J. R. Moenter S. M., Woodfikk C. I., and Karach F. J (1991). Role of thyroid gland in seasonal reproduction. Thyroxine allows a season-specific suppression of gonadotropin secretion in sheep. *Endocrinology* 129, 176-183

خالد بن أحمد البوساده

قسم وظائف الأعضاء والكيمياء الحيوية والأقربازين - كلية الطب البيطري والثروة الحيوانية
جامعة الملك فيصل - الأحساء - المملكة العربية السعودية

الملخص :

لقد تم دراسة تأثيرات الكاربازول المخصص لنشاط الغدة الدرقية في الأغنام خلال فصل الصيف. ولقد تبين أن الجرعة اليومية بمقدار ٣ ملجرام للكيلوجرام من الكاربمازول ولمدة ١١ يوماً ثبّطت عملية التبويض وتكوين الجسم الأصفر. وتشير النتائج إلى درجة الحرارة العالية والتي عادة ما تقلل من هرمونات الغدة الدرقية ربما أثرت على خصوبة الحيوانات عبر هرمونات الغدة الدرقية كما هو الحال في المعالجة بالكاربمازول.